

Animal welfare, etológia és tartástechnológia



Animal welfare, ethology and housing systems

Volume 16 Issue 1

Gödöllő
2020

SLOVAK DAIRY SHEEP – NEW COMPOSITE SHEEP BREED RAISED IN SLOVAKIA

^{1,2}Margetín Milan, ¹Janíček Martin, ¹Vavrišinová Klára, ²Pavlík Ivan, ²Huba Ján, ²Oravcová Marta, ¹Hozáková Katarína

¹Slovak Agricultural University in Nitra, Faculty of Agrobiology and Food Resources, Department of animal husbandry, 94976 Nitra (SK), Tr. A. Hlinku 2, ²NAFC–Research Institute for Animal Production Nitra, 95141 Lužianky (SK), Hlohovecká 2
Milan.Margetin@uniag.sk

Received – Érkezett: 15. 11. 2019.
Accepted – Elfogadva: 13. 05. 2020.

Abstract

In the beginning of 1990–ies, the programme of formation of synthetic population of Slovak Dairy sheep (SD) was launched. The establishment of breed was divided into two periods: first, crosses of local breeds (Improved Valachian - IV, Tsigai - Ts and Merino) with improving breeds (Lacaune - LC, East Friesian - EF) of various proportions were formed; second, a close (inter se) breeding scheme within the crossbred population was applied. At present, five thousands SD ewes are included in performance testing. Milk and reproduction traits were analyzed using mixed model methodology during the period between 1995 and 2015. The standardized milk yield (SMY) and average daily milk yield (ADMY) increased more than twice: 79.8 ± 5.36 l (1995) vs. 164.0 ± 2.04 l (2015) and 495.1 ± 33.5 ml (1995) vs. 1035.3 ± 12.8 ml (2015), respectively. In ten percent of most productive ewes, SMY was equal to 257.6 l, ADMY was equal to 1604 ml and dry matter was equal to 26.6 kg. Ewes of SD have good udder traits. Litter size in top 50 % flocks is above 150 %. The population of SD was recognized as a new dairy breed in 2017. In 2017-2019 the best flocks achieved milk production higher than 220 liters. Based on a comprehensive analysis of the genetic structure of SD we found that the overall genetic proportion of improving breeds is in new breed 60%. The remaining 40% are domestic breeds, of which IV and Ts genes are represented in the SD breed at approximately the same level (18%).

Key words: dairy ewes, composite breed, milk traits, reproduction, genetic structure

STATE OF THE ART

Following breeder's goals oriented towards high production, reproduction and functional traits of dairy sheep in Slovakia, the improving programme with an intention of forming improved sheep population fitting semi-extensive production system was launched in Slovakia in 1990–ies. This was programme of forming Slovak Dairy Sheep (SD). The breeding scheme was divided into two periods (*Margetín and Čapistrák 1994; Margetín et al., 2000a,b,c*). First, crosses of local breeds (Improved Valachian (IV), Tsigai (TS) and Merino (M)) with improving specialized breeds (Lacaune (LC), East Friesian (IF)) of various proportions were formed in some flocks involved in performance testing. Preliminary evaluations of crossbreds with proportion of LC and EF ranged from 12.5 to 87.5 % were aimed at analyses of milk traits and litter size that are considered as economically most important traits (*Apolen et al., 2000; Čapistrák et al. 2000, 2002, 2005, Margetín et al., 1993, 1999a,b*). Further analyses showed that population with good shape udder, of appropriate cistern size and

milkability was formed (*Čapistrák et al.*, 2006a,b; *Mačuhová et al.*, 2008; 2009; *Margetín et al.*, 2005b,c; *Margetín et al.*, 2011a,b; *Milerski et al.*, 2005, 2006; *Tančin et al.*, 2011). Crossbreds with various proportions of LC and IF were also confirmed to be of good growth traits both till and after weaning (*Margetín et al.* (2004a,b)). In addition, genetic parameters and variance components of udder morphology and milkability traits were preliminary studied (*Margetín et al.* 2005a, 2008).

Second, a close (inter se) breeding scheme within the crossbred population was proposed (*Margetín et al.*, 2010, 2011c, 2012b) and applied taking into account results of analyses during first period of programme. The aim was to stabilize composite population using inter se (i.e. purebred) mating scheme that included dam and sire animals used as parents of the next generation that were selected according to their breeding values for milk and litter size traits. During this period, a lot of analyses were also done. Performance testing data of flocks involved in the programme were used for evaluation. These included analyses of udder morphology and milkability traits since one goal was to form population of ewes that fit machine milking (*Margetín et al.*, 2013b; *Makovický et al.*, 2019a), have good udder morphology and functional traits (*Margetín et al.*, 2012a; *Makovický et al.*, 2013, 2014, 2015a, 2017a), are of appropriate cistern size (*Margetín et al.*, 2011d; *Makovický et al.*, 2015b,c; 2019b) and of minimal mastitis occurrence (*Margetín et al.* 2013a). Growth curves of lambs of various proportions of LC and EF within IV and TS genotypes were compared (*Makovický et al.*, 2017b). Genetic parameters of milkability traits and somatic cell score were also studied (*Makovický et al.* 2018).

In 2016, a complex analysis of SD synthetic population involving period 1995 to 2015 was done. Mixed model methodology (SAS, ver. 9.2) was employed. When milk yield during milking period (MY), standardized (SMY) and daily milk yield (DMY) as well as fat % (F%), protein % (P%), dry matter % (DM%), free of fat dry matter (DM-F%) and usable dry matter (UDM; kg) were investigated, following fixed effects: flock (52 levels), year (21 levels), parity (3 levels: first, second and third+ parity), litter size (4 levels: one lamb, two, three + lambs born and missing information about number of lambs born), covariates: milking period (in days) and interval between lambing and first test-day measurement (days) and random effect of ewe were considered. A total, 20511 records that belonged to 11026 ewes (1.86 records per ewe) entered the model for analyses of MY, SMY, DMY and 18081 records entered the model for analyses of F%, P%, DM-F% and UDM. When litter size was investigated, the following fixed effects: flock (26 levels), year (19 levels) and age of dam (8 levels, one year old up to eight+ years old) and random effect of ewe were considered. A total, 30034 litter size records that belonged to 9671 ewes entered the model.

Study of *Margetín* (2016) confirmed that improving programme that resulted in acknowledgement of SD sheep with more favourable traits than those of local breeds was fulfilled. During period of 21 years (from 1995 to 2015), SMY increased about twice (from 79.8 l in 1995 to 167.0 l in 2015). Similarly, DMY and production of usable dry matter increased by 109 and 79 %. Most productive flocks reached MY ranging from 170 to 190 l. Top 10 % ewes reached SMY above 250 l, DMY reached more than 1600 ml and production of usable dry matter was above 26 kg. Average F% and P% slightly decreased (7 and 2.4 %, respectively). Also, ewes of SD population were of good udder morphology and functional udder traits. Prolificacy did not fulfill expectations and was equal to 141.1 %. Nevertheless, it was about 160 % in top flocks. Average daily gain till weaning was about 300 g in male lambs and about 260 g in female lambs. The frequency of ARR allele of the prion gene ranged from 0.711 to 0.789 (*Margetín et al.*, 2016).

Results of SD sheep flocks resulted in fact that this population was recognized as a native breed on May 26, 2017. Detailed information about history of breeding this population can be found in study of *Margetín et al.* (2017). Moreover, the importance of development of

this population from social and breeder's point of view is described in study of Margetín (2017a,b). In 2017-2019 the best flocks achieved milk production higher than 220 liters. The average fat content of ewes in the performance control (15 flocks) was 7.01 resp. 7.03 % and protein content of 5.60 resp. 5.43% in 2017 and 2018. Based on a comprehensive analysis of the genetic structure of SD we found that the overall genetic proportion of improving breeds (LC, EF) is in new breed 60%. The remaining 40% are domestic breeds, of which IV and Ts genes are represented in the SD breed at approximately the same level (18%; Margetín et al. (2018b)). Based on the analysis of SD breed, we can conclude that the genes of LC breed are more prevalent (54 %) in the SD breed than the EF genes (6 %), although the chances of both improving breeds at the beginning of the breeding process were equal (breeding programs recommended use both breeds).

CONCLUSION

Population of SD breed is accordingly large (more than 5 ths. females in performance testing, 35 % yearling females enter mating schemes), so it can be successfully bred in Slovakia in the future. Breeders are satisfied with level of production and reproduction traits reached (Pavlik et al., 2017; Chrenek et al., 2018; Margetín et al., 2018a, 2019). With respect to appropriate effective population size, it is important that more than 300 sires of Slovak Dairy sheep are produced each year. In the next years, not only numbers of SD population, its production and reproduction traits, but also its exterior traits including wool characteristics need to be continuously stabilized (Margetín, 2017c).

ACKNOWLEDGEMENTS

The study was supported by the projects VEGA 1/0364/15 and KEGA 015SPU-4/2019.

REFERENCES

- Apolen, D., Čapistrák, A., Margetín, M. (2000): Mlieková úžitkovosť a plodnosť kríženiek plemena merino s dojnymi plemenami. Acta fytotechnica et zootechnica. Nitra : XIX. Dni genetiky, 3. 121.
- Čapistrák, A., Margetín, M., Apolen, D., Špánik, J. (2000): Produkcia mlieka a plodnosť oviec plemena zošľachtená valaška, kríženiek s plemenami lacaune a východosfrízskym plemenom. Acta fytotechnica et zootechnica. Nitra : XIX. Dni genetiky, 3. 120.
- Čapistrák, A., Margetín, M., Apolen, D., Špánik, J. (2002): Produkcia a obsah základných zložiek mlieka oviec plemena zošľachtená valaška, lacaune a ich kríženiek. Vedecké práce VÚŽV Nitra, 35. 89–96.
- Čapistrák, A., Margetín, M., Apolen, D., Špánik, J. (2005): Produkcia a zloženie mlieka bahníc rôznych plemien a ich kríženiek počas dojnej períody. Vedecké práce VÚŽV Nitra, 38. 181–189.
- Čapistrák, A., Margetín, M., Apolen, D., Špánik, J. (2006a): Porovnanie morfologických ukazovateľov vemena rôznych genotypov oviec. Acta fytotechnica et zootechnica, 9. mimoriadne vydanie. 179–181.
- Čapistrák, A., Margetín, M., Špánik, J., Apolen, D. (2006b): Lineárne hodnotenie, miery vemena a produkcia mlieka rôznych genotypov oviec. In: Zb. prednášek „Ovce , Kozy Seč 2006“, 10.-11.11.2006. 106-113. ISSN 1213-600X.
- Chrenek, P., Margetín, M., Bulla, J., Kubovičová, E., Kulikova, B., Makarevich, A. (2018): New Slovak animal breeds. In *Danubian Animal Genetic Resources*. Proceedigs of 29th Annual Meeting of Dagene. 24.-27.6.2018. Hungary, 62-64. ISSN 2498-5910.

- Mačuhová, L., Uhrinčat', M., Mačuhová, J., Margetín, M., Tančin, V. (2008): The first observation of milkability of the sheep breeds Tsigai, Improved Valachian and their crosses with Lacaune. Czech Journal of Animal Science, 53. 12. 528–536.
- Mačuhová, L., Uhrinčat', M., Mačuhová, J., Tančin, V. (2009): Milkability of Tsigai, Improved Valachian, and their crosses with Lacaune. Acta fytotechnica et zootechnica, 12. spec. iss. 385–394.
- Makovický, P., Nagy, M., Makovický, P. (2013): Comparison of external udder measurements of the sheep breeds Improved Valachian, Tsigai, Lacaune and their crosses. Chilean Journal of Agricultural Research, 73. 4. 366–371.
- Makovický, P., Nagy, M., Makovický, P. (2014): The comparison of ewe udder morphology traits of Improved Valachian, Tsigai, Lacaune breeds and their crosses. Mljetkarstvo, 64. 2. 86–93.
- Makovický, P., Margetín, M., Makovický, P. (2015a): Genetic parameters for the linear udder traits of nine dairy ewes , short commuinication. Veterinarski Archiv, 85. 5. 577–582.
- Makovický, P., Margetín, M., Milerski, M. (2015b): Estimation of udder cistern size in dairy ewes by ultrasonography. Mljetkarstvo, 65. 3. 210–218.
- Makovický, P., Milerski, M., Margetín, M., Makovický, P., Nagy, M. (2015c): Genetic parameters for the size of udder cisterns in ewes diagnosed by ultrasonography among breeds: Improved Valachian, Tsigai, Lacaune and their crosses. Archivos de zootecnia, 64. 248. 403–408.
- Makovický, P., Margetín, M., Makovický, P. (2017a): Estimation of genetic and phenotypic parameters for udder morphology traits in different dairy sheep genotypes. Acta Universitatis Agriculturae et Silviculturae Mendelianae Brunensis, 65. 1. 105–110.
- Makovický, P., Margetín, M., Nagy, M., Makovický, P. (2017b): Growth curves in lambs of various genotypes created on the basis of Improved Valachian and Tsigai breeds. Acta Universitatis Agriculturae et Silviculturae Mendelianae Brunensis, 65. 1. 111–118.
- Makovický, P., Margetín, M., Nagy, M., Makovický, P. (2018): Genetic parameters for milkability and somatic cell score in Slovak dairy sheep. Indian Journal of Animal Sciences, 88. 7. 860–862.
- Makovický, P., Margetín, M., Makovický, P., Nagy, M. (2019a): Machine milkability of East Friesian and Lacaune dairy sheep. Indian Journal of Animal Sciences, 89. 6. 686–691.
- Makovický, P., Nagy, M., Margetín, M., Poráčová, J., Milerski, M., Makovický, P. (2019b): Measurement of the mammary gland cistern of dairy ewes. Acta Universitatis Agriculturae et Silviculturae Mendelianae Brunensis, 67. 4. 957–961.
- Margetín, M. (2016): Aktuálny stav šľachtenia slovenskej dojnej ovce. Chov oviec a kôz, 36. 4. 27–37.
- Margetín, M. (2017a): Nové národné plemeno oviec - slovenská dojná ovca. Poľnohospodár, 61. 10. 2.
- Margetín, M. (2017b): Slovenská dojná ovca , praktický výsledok inovačného procesu. Agromagazín, 19. 7. 36–37.
- Margetín, M. (2017c): Slovenská dojná ovca , ako ďalej v šľachtení. Chov oviec a kôz, 37. 2. 3–4.
- Margetín, M., Čapistrák, A., Kališ, M. (1993): Produkcia mlieka cigájskych bahníc a kríženiek s východofrízskym plemenom vo vzťahu k hmotnosti jahniat v odchovnej fáze. Journal of Farm Animal Science, 26. 181–188.
- Margetín, M., Čapistrák, A. (1994): Zámery v šľachtiteľskom programe chovu oviec, kontrole úžitkovosti a dedičnosti. In: *Predpoklady efektívneho chovu oviec v súčasnosti*. Trenčín : Výskumný ústav živočíšnej výroby, Stanica chovu a šľachtenia oviec a kôz, 8–19.
- Margetín, M., Apolen, D., Čapistrák, A. (1999a): Zošľachtovanie merinských oviec v smere mäsovo , mliekovom. Nás chov, 59. 5. 17–19.

- Margetín, M., Čapistrák, A., Špánik, J., Apolen, D., Hyžová, A. (1999b): Produkcia, zloženie mlieka a laktačné krvky bahníc plemena zošlachtená valaška, lacaune a ich kríženiek v období cicania jahniat. *Journal of Farm Animal Science*, 32. 111–119.
- Margetín, M., Čapistrák, A., Apolen, D. (2000a): Program zošlachťovania merinských oviec s použitím plemena lacaune a východofrízskeho plemena v šlachtitel'sko - experimentálnych chovoch. *Chov oviec a kôz*, 21. 1,2. 24–27.
- Margetín, M., Čapistrák, A., Apolen, D. (2000b): Program zošlachťovania cigájskych oviec s použitím plemena lacaune a východofrízskeho plemena v šlachtitel'sko - experimentálnych chovoch. *Chov oviec a kôz*, 21. 1,2. 27–30.
- Margetín, M., Čapistrák, A., Apolen, D. (2000c): Program zošlachťovania valašských oviec s použitím plemena lacaune a východofrízskeho plemena v šlachtitel'sko - experimentálnych chovoch. *Chov oviec a kôz*, 21. 3. 4–7.
- Margetín, M., Bullová, M., Čapistrák, A. (2004a): Rastové krvky jahniat rôznych genotypov vytvorených na báze plemena zošlachtená valaška a cigája. In: *Aktuálne problémy riešené v agrokomplexe [CD]*. Nitra : Slovenská poľnohospodárska univerzita v Nitre, 456-461. ISBN 80-8069-488-6.
- Margetín, M., Čapistrák, A., Špánik, J., Apolen, D., Bullová, M. (2004b): Rastová intenzita jahniat pri tvorbe rôznych genotypov plemena cigája. In: *Chov hospodárskych zvierat v podmienkach EÚ [CD]*. Nitra : Slovenská poľnohospodárska univerzita v Nitre, 331–337.
- Margetín, M., Čapistrák, A., Apolen, D., Oravcová, M., Špánik, J. (2005a): Doterajšie výsledky programu zošlachťovania dojných oviec na Slovensku. In: *Zborník z medzinárodnej vedeckej konferencie „Ekologické a ekonomicke aspekty využitia poľnohospodársky znevýhodnených plôch chovom malých prežívavcov“*. Žilina : VÚM, 115-122.
- Margetín, M., Milerski, M., Apolen, D., Čapistrák, A., Oravcová, M. (2005b): Morphology of udder and milkability of ewes of Tsigai, Improved Valachian, Lacaune breeds and their crosses. In: Conference on „Physiological and technical aspects of machine milking. Nitra : Slovak Republic, 26. - 28. apríl 2005, ICAR Technical series no. 10. 259-263. ISSN 1563-2504, ISBN 92-95014-07-3
- Margetín, M., Špánik, J., Milerski, M., Čapistrák, A., Apolen, D. (2005c): Relationships between morphological and functional udder traits and somatic cell count in milk of ewes. In: Conference on „Physiological and technical aspects of machine milking. Nitra : Slovak Republic, 26. - 28. apríl 2005, ICAR Technical series no. 10. 255-258. ISSN 1563-2504, ISBN 92-95014-07-3
- Margetín, M., Makovický, P., Čapistrák, A., Apolen, D., Oravcová, M., Tančin, V. (2008): Odhad genetických parametrov vybraných ukazovateľov charakterizujúcich morfológiu vemena a dojiteľnosť bahníc. In: CD , zb. z medz. ved. konferencie „Aktuálne problémy riešené v Agrokomplexe“, 5.12.2008, Nitra : SPU. 234-240. ISBN 978-80-552-0151-1
- Margetín, M., Apolen, D., Čapistrák, A., Špánik, J., Margetínová, J. (2010): Postup šlachtenia syntetickej populácie oviec vytvorennej s použitím plemena lacaune a východofrízske. In Rozvoj metód zlepšovania a efektívnejšieho využívania genofondu zvierat z hľadiska konkurencieschopnosti, kvality produktov a ich zdravotnej bezpečnosti. Nitra : Centrum výskumu živočíšnej výroby Nitra, Ústav chovu a šlachtenia oviec a kôz Trenčianska Teplá, 57 p.
- Margetín, M., Apolen, D., Čapistrák, A., Kulinová, K. (2011a): Results of improvement program in dairy sheep raised in Slovakia with using of Lacaune and East Friesian breed. In *Możliwości poprawy efektywnoci chowu owiec i kóz w Polsce w ujęciu regionalnym*. Kraków : Instytut Zootechniki.. 59. ISBN 978-83-7607-188-6.

- Margetín, M., Apolen, D., Čapistrák, A., Milerski, M., Oravcová, M., Debrecéni, O. (2011b): Vplyv špecializovaných dojných plemien na veľkosť cisterien vemena kríženiek valašských oviec. *Acta fytotechnica et zootechnica*, 13. 4. 85–92.
- Margetín, M., Čapistrák, A., Apolen, D., Reľovský, S., Šutý, J., Rafajová, M. (2011c): Tvorba syntetickej populácie slovenskej dojnej ovce, druhá etapa programu zošľachtovania. *Chov oviec a kôz*, 21. 4. 6–8.
- Margetín, M., Makovický, P., Milerski, M., Apolen, D., Debrecéni, O., Oravcová, M. (2011d): The effect of specialized dairy breeds on udder cistern size in Tsigai crossbreeds. *Slovak Journal of Animal Science*, 44. 4. 146–153.
- Margetín, M., Oravcová, M., Apolen, D., Milerski, M. (2012a): Genetic parameters for udder traits in Slovak dairy sheep and their crosses with specialized breeds. *Journal of Life Sciences*. 6. 454–459.
- Margetín, M., Apolen, D., Čapistrák, A. (2012b): Šľachtenie slovenskej dojnej ovce, druhá etapa zošľachtovacieho programu. *Slovenský chov*, 17. 1. 25–27.
- Margetín, M., Milerski, M., Apolen, D., Čapistrák, A., Oravcová, M., Debrecéni, O. (2013a): Relationships between production, quality of milk and udder health status of ewes during machine milking. *Journal of Central European Agriculture*. 14. 1. 328–340.
- Margetín, M., Oravcová, M., Makovický, P., Apolen, D., Debrecéni, O. (2013b): Milkability of Improved Valachian, Tsigai and Lacaune purebred and crossbred ewes. *Slovak Journal of Animal Science*, 46. 3. 100–109.
- Margetín, M., Bireš, J., Reľovský, S., Oravcová, M., Mojžiš, M., Rajská, Z. (2016): Changes in allele and genotype frequencies of *PrP* gene in breeds of sheep in Slovakia between 2004 and 2015. *Acta fytotechnica et zootechnica*, 19. 3. 119–122.
- Margetín, M., Apolen, D., Čapistrák, A. (2017): História procesu šľachtenia slovenskej dojnej ovce. *Slovenský chov*, 22. 5. 14–17.
- Margetín, M., Čapistrák, A., Apolen, D., Huba, J., Janíček, M. (2018a): Proces tvorby a úžitkové parametre nového národného plemena oviec – slovenská dojná ovca. In *Zborník abstraktov „Genetická konferencia GSGM“*. 12.-14.9.2018, Bratislava, 71. ISBN 978-80-229-4545-3.
- Margetín, M., Šulko, K., Reľovský, S., Oravcová, M., Janíček, M. (2018b): Zmeny genetickej štruktúry syntetickej populácie oviec v procese tvorby plemena slovenská dojná ovca. *Chov oviec a kôz*, 37. 4. 4–7.
- Margetín, M., Šutý, J., Šulko, K., Oravcová, M., Janíček, M. (2019): Výsledky šľachtenia dojných oviec na mliekovú úžitkovosť za posledné štvrtstoročie. *Slovenský chov*, 24. 1-2. 14–17.
- Milerski, M., Margetín, M., Apolen, D., Čapistrák, A., Špánik, J. (2005): Udder cistern size and milkability of ewes of various genotypes. In: Conference on „Physiological and technical aspects of machine milking. Nitra : Slovak Republic, 26.-28. apríl 2005, ICAR Technical series No. 10. 63–69.
- Milerski, M., Margetín, M., Čapistrák, A., Apolen, D., Špánik, J., Oravcová, M. (2006): Relationships between external udder measurements and the linear scores for udder morphology traits in dairy sheep. *Czech Journal of Animal Science*, 51. 9. 383–390.
- Pavlík, I., Margetín, M., Janíček, M. (2017): Slovenská dojná ovca , výsledky u popredných chovateľov sú príslušom budúcnosti. *Chov oviec a kôz*, 37. 2. 17–18.
- Tančin, V., Mačuhová, L., Oravcová, M., Uhrinčat', M., Kulinová, K., Roychoudhury, S., Marnet, P. G. (2011): Milkability assessment of Tsigai, Improved Valachian, Lacaune and F1Crossbred ewes (Tsigai x Lacaune, Improved Valachian x Lacaune) throughout lactation. *Small Ruminant Research*, 97. 28–34.