

Camera traps as a research method for carnivore population estimation: Strength, weaknesses, opportunities and threats, analysis and improvements

Alexandra KALANDARISHVILI^{1*} – Miklós HELTAI¹

1: Department of Wildlife Biology and Management, Institute for Wildlife Management and Nature Conservation, Hungarian University of Agriculture and Life Sciences, Gödöllő, Páter Károly u. 1, 2100 Hungary., e-mail: alexkalendarishvili97@gmail.com

Abstract: Camera traps have been gaining popularity in population estimation studies. Based on 60 scientific journals, we evaluated the strengths, weaknesses and improvements of the camera trap method to better understand its effectiveness for studying population parameters. Camera traps have a strong advantage of being a non-invasive method, requiring minimal labor and because of its ability to detect multiple species per sampling effort. However, theft and time-consuming data analyses, poor sensor performance and potential behavioral changes of wildlife due to noise and flashlights, prevent the camera traps from being the optimal population estimation method. The population parameter studied depends strongly on the behavior and biology of the target species, although the most common opportunity for development is all related to sensor performance (better triggering response and higher sensitivity) as well as extreme weather condition resistance.

Keywords: Monitoring, Research methods, Large carnivores, Method analysis

Received 24 July 2023, Revised 19 December 2023, Accepted 20 December 2023

 This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License

Introduction

Estimating carnivore population abundance is an active challenge, nevertheless, obtaining such information as accurately as possible is critical (Nichols & Williams, 2006). Camera trapping has emerged as a powerful tool that allows for non-invasive data collection to study behavioral and ecological aspects of target species (Delisle et al., 2021). Information from camera traps allow unambiguous individual identification making its data useful for generating accurate population estimates from capture-recapture analysis (Joubert et al., 2020; Macdonald et al., 2020).

Camera traps are being actively used to study carnivore population diversity and evaluate their abundance and in recent years cameras

have been used to study more complex elements of carnivore populations such as age structures, predatory behavior, and daily activities (Joubert et al., 2020; Miyamoto et al., 2018; Thornton et al., 2018). However, there are several factors that may impact the overall performance of the camera traps: users' expertise, the condition of the study area, the quality of the equipment, and bias due to systematic error. Certain limitations – technical, user or otherwise are further limited by great differences among the cameras themselves, namely in their sensitivity, detection zones and performance under variable environmental conditions.

Comparative tests of the applicability and effectiveness of camera traps are rarely conducted, although it is important to know the potential issues in terms of the strengths and

weaknesses and select traps suitable for local application. In this paper, we aim to conduct a systematic literature review and analyze camera trapping as the method to estimate wildlife populations using a known method of SWOT analysis (strength, weaknesses, opportunities, and threats).

Materials and Methods

We used the Scopus as our database to gather publications. The keywords that we combined included “camera trap” OR “photo trap” OR “remote camera” AND “effectiveness” OR “Practicality” AND “Carnivores” OR “Predators”. The search for scientific papers was filtered by keywords which were included in the title, abstract and/or keywords. To perform the SWOT analysis, we used the following definitions to extract information from the scientific papers and classify them accordingly. *Strength* in this case is defined as attributes of the camera trap methodology that benefits the study at hand and makes it easier and more practical to execute. In other words, characteristics that are absent in other more traditional methods and that separate camera trapping from other methods. *Weaknesses* are defined by attributes that hinder the method to perform at its optimal level and attributes that are likely to lead to biased or imprecise results. We defined the *Opportunities* as potential external factors that are not part of the method but could potentially give a method an additional advantage. Finally, we defined *Threats* as factors that have the potential to harm the final outcome of the study.

Results

In total 60 articles were generated as a result of the search. The list of the scientific publications that were collected does not include all papers that were published on the

topic of camera trapping, nevertheless, the list acquired enables us to have a good overall understanding of the limitations and advantages of the camera traps as a population estimation method. The strengths, weaknesses, opportunities and threats of the camera trap method based on the analyzed papers are summarized in Table 1.

Discussion

Strengths and Weaknesses

We investigated the strengths and weaknesses of camera trapping as a first part of the study. Camera trapping is relatively low cost in the long term compared to other methods of population estimation, it is non-invasive, and makes it possible to obtain information on trap-shy species in a wide range of habitats (Table 1 Strength 1 & 5). It is a great tool that replaces traditional methods especially in areas that are remote and hard to access (Table 1 Strength 4) while producing data on multiple species simultaneously (Table 1 Strength 6). Steinbeiser et al. (2019) demonstrated the substantial underestimation of species richness using transect surveys in comparison to camera traps in a savanna ecosystem.

Recorded photos permanently document multiple types of information. Camera-traps are now being more frequently used to study behaviors of species such as unique behavioral associations among and between different trophic levels (Table 1 Strength 7). For example, Burton et al. (2012) modeled the responses of carnivores to hunting, habitat and prey in western African protected areas, while Thornton et al. (2018) managed to document two spatial hotspots of probable hunting association between badgers and coyotes in north-central Washington which suggests a large role of environmental characteristics in shaping foraging associations.

Camera traps are specifically effective methods for mid-sized and large carnivore species

Table 1: SWOT analysis of the camera trapping as a method for population estimation.

Strengths
<ol style="list-style-type: none">1. Not invasive method of data collection (Gompper et al., 2006; Joubert et al., 2020)2. Effective monitoring method for mid-sized and large carnivore species (Avrin et al., 2021; Balme, Slotow, & Hunter, 2009; Gompper et al., 2006; Joubert et al., 2020; Kämmerle et al., 2019; Rogan et al., 2022; Shamoan et al., 2017; Strampelli et al., 2020; Stobo-Wilson et al., 2020; Sunarto et al., 2015)3. Most optimal choice for the the studies focusing on population diversity and abundance evaluation (Balme, Slotow, & Hunter, 2009; Beirne et al., 2021; Burgar et al., 2019; Farris et al., 2014; Kluever et al., 2013; Lazenby et al., 2015; Macdonald et al., 2020; Muench & Martínez-Ramos, 2016; O'Brien & Kinnaird, 2011; Palencia et al., 2021; Rogan et al., 2022; Shamoan et al., 2017; Silveira et al., 2003; Steinbeiser et al., 2019; Xiao et al., 2016)4. Great replacement for traditional methods when working in remote, hard to access areas (Bernard et al., 2022; Silveira et al., 2003; Steinbeiser et al., 2019)5. Minimal labor and cost effective on a long term (Delisle et al., 2021; Foresman & Pearson, 1998; Steinbeiser et al., 2019)6. Ability to detect multiple species and Species detection is usually unambiguous (Burgar et al., 2019; Farris et al., 2014; Foresman & Pearson, 1998; Joubert et al., 2020; Macdonald et al., 2020; O'Connell et al., 2011)7. Recorded photos are permanent and document multiple types of information (age, behavior, predation, daily activity) (Comer et al., 2018; Joubert et al., 2020; Lazenby et al., 2015; Miyamoto et al., 2018; O'Connell et al., 2011; Srbek-Araujo et al., 2017; Thornton et al., 2018)
Weaknesses
<ol style="list-style-type: none">1. Behavior changes of wildlife (Meek et al., 2016; Selonen et al., 2022)2. Ineffective for small carnivores (Gompper et al., 2006; Pirie et al., 2016)3. Needs to be accompanied with other methods for accuracy (Balme, Hunter, & Slotow, 2009; Engeman & Witmer, 2000; Gompper et al., 2006; Pirie et al., 2016)4. Time-consuming data analyses and expensive equipment (Foresman & Pearson, 1998; Glover-Kapfer et al., 2019; Kelly et al., 2012; O'Connell et al., 2011; Palencia et al., 2021; Steinbeiser et al., 2019)5. Battery and memory limitations (O'Connell et al., 2011)6. High risk of theft and wildlife damage (Glover-Kapfer et al., 2019; Kelly et al., 2012; Steinbeiser et al., 2019)7. Camera traps have limitations for estimating population parameters, especially if individuals cannot be identified (Jordan et al., 2011; Joubert et al., 2020; Kelly et al., 2012; Larrucea, Brussard, et al., 2007; Larrucea, Serra, et al., 2007; Negrões et al., 2010)

Table 1: SWOT analysis of the camera trapping as a method for population estimation (continued).

Opportunities
<ol style="list-style-type: none"> 1. Allows for complex sampling design for studies on a population level (Burton et al., 2012; Farris et al., 2014; Kämmerle et al., 2019; Kelly, 2001; Kluever et al., 2013; Mendoza et al., 2011; Rogan et al., 2022; Sunarto et al., 2015; Thornton et al., 2018) 2. Can be used with or without bait/lure (Barcelos et al., 2023; Buyaskas et al., 2020; Comer et al., 2018; Joubert et al., 2020; Dri et al., 2022; Palmer et al., 2021; Stobo-Wilson et al., 2020; Thorn et al., 2009) 3. Can be paired with other software and give more diverse results (Mendoza et al., 2011) 4. Camera traps are used to generate abundance indices (Beirne et al., 2021; Burton et al., 2012; Farris et al., 2014; Muench & Martínez-Ramos, 2016; Sollmann et al., 2013; Strampelli et al., 2020; Windell et al., 2022) 5. Pictures produced are aesthetically pleasing and can be used in research fundraising and awareness raising (O’Connell et al., 2011) 6. Can be used to evaluate other monitoring measures (Avrin et al., 2021; Comer et al., 2018; Nekaris et al., 2020; Palmer et al., 2021; Windell et al., 2022)
Threats
<ol style="list-style-type: none"> 1. Capturing non-target species (Kelly et al., 2012) 2. Human scent may cause the animals to avert the camera device (Buyaskas et al., 2020) 3. Bias due to under-recording, misidentification of the species and/or translating the data (Borah et al., 2014; Joubert et al., 2020; Mendoza et al., 2011; Pirie et al., 2016) 4. Data can be lost due to equipment failure (O’Connell et al., 2011) 5. Technology is changing rapidly and software needs to be constantly updated (Mendoza et al., 2011)

(Table 1 Strength 2). While some authors like Gompper et al. (2006) and Pirie et al. (2016) argue that camera traps are ineffective for observe and study small mammal, Srbek-Araujo et al. (2017) successfully investigated squirrel preys on seeds defecated by lowland tapirs in the Atlantic Forest, southeastern Brazil.

There are two parameters to assess the effectiveness of a method: latency to initial detection (LTD) and probability of detection (POD). LTD measures the time it takes for the first detection of a species at a survey site to be documented. POD looks at the probability of detecting a species with a specific

technique. In an ideal scenario, more efficient survey methodologies should result in a low LTD and high POD.

Gompper et al. (2006) found that the value for each of those parameters varies depending on the target species and concluded that for mid-sized carnivores, for example, raccoon, fisher, opossum, and domestic cat, camera traps and plate tracks had a very similar detection efficiency, this is also confirmed by the study done by Shamoan et al. (2017) that investigated medium-sized carnivores in Mediterranean agricultural areas. At the same time, the wariness and aversion to foreign equipment by the animals showed

higher LTD for track plates compared to camera traps (Gompper et al., 2006).

LTD can be manipulated through a combination of bait and lure on the site (Table 1 Opportunities 2). Buyaskas et al. (2020) finds that the combination of bait and lure as an attractant was particularly effective for all mustelid species, especially American marten and fisher, and slightly less effective than bait for short-tailed weasel. Opossum records in lured stations were almost three times higher than in non-lured stations Barcelos et al. (2023). Lure has proved comparatively more effective than bait for American black bear and bait was notably more effective than lure for mustelids, which also had a much greater chance of being detected with attractive use than other carnivores (Buyaskas et al., 2020).

For smaller mammals like martens and weasels, track plates had higher POD compared to camera traps, however, camera traps proved to be a useful method to survey the bears showing low LTD and high POD (Gompper et al., 2006). On the other hand, because coyotes are more wary, cameras proved ineffective as shown by the high LTD and low POD. For coyotes, the best method for surveying remains to be snow tracking (Gompper et al., 2006). Buyaskas et al. (2020) also confirms this by concluding that compared to mustelids, the use of attractants for eastern coyote and American black bear was less successful in maximizing detection probability, despite increases in detection probability for both species, suggesting that the eastern coyote is wary of human scent at bait stations.

Buyaskas et al. (2020) found that except for coyotes and red foxes, the faecal surveys proved inefficient to detect the presence of the species. Genetic tests of the fecal and snow tracking confirmed the presence of red foxes in areas where other methods were unable to document them. As a result, Gompper et al. (2006) argue that cam-

eras and track-plates are inefficient for surveying small canids in some harder-reaching regions. The high POD indicated that snow tracking surveys were highly effective for detecting species that are normally active in winter, and this method may be more effective than both cameras and track plates given that the conditions are suitable.

Behavior changes of wildlife as a result of camera traps was also observed by Meek et al. (2016) and Selonen et al. (2022) who argue that the flash light causes animals to avoid the camera stations (Table 1 Weakness 1). Dealing with these changed reactions is critical as this will influence time spent in front of the camera, and can therefore result in bias. As a result, some authors proposed to disregard the first period of the survey to allow animals to become used to the equipment (Howe et al., 2017); while others discarded all the observations where animal behavior indicated a slight change in reaction to the camera traps (Bessone et al., 2020). while others discarded some data to obtain a reasonable detection function fitting (Cappelle et al., 2019). This all points to the fact that different methods need to be used when trying to examine different species in the carnivore community.

Those studies that set out to report on wildlife species on a population or community level require a much more intense sampling effort and a more complex design of the sampling technique. Camera traps have been actively used in that regard, as they are said to offer a better alternative to the so-called traditional methods that focus on population diversity and abundance evaluation (Table 1 Strength 3). However, camera traps have limitations for estimating population parameters especially if individuals cannot be individually identified and this is only possible for species with distinctive markings, (Table 1 Weakness 7).

Moreover, analyzing the data from the camera traps is extremely time consuming

and requires purchasing of rather expensive equipment (Table 1 Weakness 4) which are in themselves limited by battery life and memory, thus in need constant monitoring (Table 1 Weakness 5). Cameras being an expensive and a valuable tool are often at risk of theft or damage by wildlife resulting in both financial and experimental losses (Table 1 Weakness 6), although this is easy to prevent by use of locks, camouflage, or security cases.

Nevertheless, Silveira et al. (2003) mentions that when it comes to comparing different methods and choosing one for the population diversity and abundance evaluation, camera traps are the better choice, primarily because camera traps are most useful and appropriate in remote areas that are difficult to access and where conducting traditional methods like line transects and/or animal track/scat surveys are rather impossible (Table 1 Strength 4). Contrary to Silveira et al. (2003), Gompper et al. (2006) and Pirie et al. (2016) argue that camera traps are quite incapable to identify small canids that would otherwise be easily detected by more traditional methods for example by scat or track surveys and/or DNA analysis (Gompper et al., 2006; Silveira et al., 2003). Interestingly, Pirie et al. (2016) similarly found that in South Africa camera traps largely under-recorded the number of animals that were passing a trapping area compared to those identified using traditional traps, especially the smaller species. The study effectively demonstrated that the track plates can provide us with an opportunity to advance the success of the camera traps. In order to avoid under-recording of the species Engeman and Witmer (2000); Gompper et al. (2006); Balme, Hunter, and Slotow (2009); Pirie et al. (2016) argue that camera trapping stations need to be accompanied by other methods to improve accuracy of species recording (Table 1 Weakness 3).

Threats and Opportunities

It is true that camera traps allow for com-

plex sampling design for studies that focus on population level. For example, using camera traps as the main method, Sunarto et al. (2015) successfully addressed knowledge gaps on the topic of cat coexistence in central Sumatra by investigating general ecological characteristics of each cat species in relation to geographic location and site conditions; factors affecting probability of site use by each cat species; and the extent of interactions between cat species pairs as indicated by spatial and temporal co-occurrence. However, it is important to consider the bias of the camera trapping for the population estimation studies and account for that bias. Bias due to under-recording, misidentification of the species and/or translating the data (Table 1 Threats 3).

Consideration of human presence at the monitoring site should also be taken into account in the initial study design. If the study design protocol requires frequent site visits for rebating as concluded by Barcelos et al. (2023), care must be taken to assure that the species of study is known to be resilient to effects of human presence; otherwise, such a protocol may not be suitable as human scent may cause the animals to avert the camera device (Buyaskas et al., 2020). Lure renewing will in itself imply a significant increase in field-related costs and it's likely to bias other species studies.

Camera traps are used to generate abundance indices as well, to get a quick insight into population size (Table 1 Opportunities 4). However, it is important to note that the indices are in themselves limited and biased compared to actual population density. It is important to note that differences and variations in indices are not directly proportionate to variations in the actual population size. However, the very nature of using the indices requires the researcher to establish certain assumptions. For example, one such assumption is that wildlife detectability is constant in both dimensions of space and time as

well as among species which in itself is questionable (Sollmann et al., 2013). Moreover, indices are not often corrected to the actual population dynamics, this leads to the indices being unable to give insight into the true population dynamics (Sollmann et al., 2013).

Nonetheless, the information gathered from camera traps can prove a great use in the occupancy model which aims to study species occurrence and absence from the area in order to outline certain population dynamic parameters. This has very helpful implications for monitoring elusive species for which observations are scarce (Trollet et al., 2014). Once the information is gathered from the cameras, researchers must realize that data processing often takes more time than deploying and monitoring cameras in the field (Table 1 Weakness 4). Considerable data management is required in any camera-trapping study such as sifting through photographs and entering them into a relevant database. In addition, other than target species, cameras also capture nontarget species which also need to be entered into the database (Kelly et al., 2012). While it may seem like nontarget species are an unimportant part of the study at hand, they can also provide some important information, for example, nontarget species can give a great idea about potential competitors, and the distribution of prey. It is also a possibility to link the trapping rates of the target carnivores to the trapping rates of prey (Kelly et al., 2012).

Individual animals are identified by natural fur marks, injuries, and coloration patterns. While the differences in the individual markings may be obvious, these types of identifications are always subjective and will vary depending on the individual observer. This also affects the precision of the estimation outcome. In order to minimize the possibility to misidentify an individual, a number of computer models have been developed for this specific reason - to help identify the pictures of marked animals (Kelly, 2001; Men-

doza et al., 2011). These tools significantly improve the researchers' ability to recognize and identify individuals and ultimately make population density estimates more accurate.

Improvements

In their review of 2,167 papers, Delisle et al. (2021) observed that there is a significant decline in studies published since 2005 that used capture-recapture camera trap methodology as the main study method. One of the most common reasons for these declines in camera trap use is researchers' inability to effectively identify individuals. As a result, many researchers who are unable to identify individuals either shifted their focus to estimates of occupancy or switched to using abundance indexes. Both alternatives are less costly choices (Delisle et al., 2021).

To tackle the issue of identification, Joubert et al. (2020) designed an study site arrangement by strategically positioned baits to enhance the identification of individually marked carnivores, ensuring optimal scrutiny of the right side for species with distinct coat patterns like leopards, jaguars, ocelots, and clouded leopards. This approach facilitates precise measurement of body dimensions, as well as more accurate determination of an individual's age and sex from photographs. However, a potential limitation lies in the methodology favoring the recording of individuals attracted to bait, potentially introducing bias by not capturing bait-shy individuals. Moreover, the use of baits at camera stations may influence the ranging behavior of the target species, potentially conditioning them to bait presence and affecting their movement patterns during sampling.

In order to have more efficient camera trap data that leads to unbiased population estimations, it is important to consider the distribution of cameras around the study site depending on the habitat use, compliance with the assumptions of mark-recapture models, sampling frequency, and the adequate selection of individual animal characteristics to

be used to distinguish between different individuals (Mendoza et al., 2011). However, one other methodological aspect that needs to be given significant attention is the reduction of subjectivity during individual identification.

Mendoza et al. (2011) suggested two ways in which misidentification and its potential bias can impact population size estimates. First, in case the misidentification cannot be improved, models can be composed that would account for the misidentification and include its possible effects in the final population estimations (Mendoza et al., 2011). Yoshizaki et al. (2009) modeled the possible effect that miscounting would have on the population estimates due to natural marks on a single individual changing over time, for example, evolving markings over time contributed to a significant bias and often overestimated the population size (Yoshizaki et al., 2009). The second way to prevent misidentification bias is to digitalize the identification process. This is primarily important when a researcher works with exceptionally large picture databases which are repetitive, and time-consuming, therefore, easily susceptible to identification error. Kelly (2001), used a 3-dimensional computer-matching system to assist to classify close to 10,000 images of Serengeti cheetahs. While the speed of identification increased dramatically as well as identification accuracy, the effectiveness of the method highly depends on the quality of the image and the angle at which the compared pictures have been captured (Kelly, 2001).

These models, while theoretically viable, have two practical shortcomings when identifying species. The first shortcoming is the rather poor performance of the camera traps for rare species, and the second shortcoming is the overall poor transferability. This is typical for the case scenarios when classifying pictures from the cameras that are not in the model training set (Delisle et al., 2021). Poor

detection of rare species is a cause for actual concern as the rare species are of greater conservation interest. However, the poor transferability is a significantly bigger issue as it in turn limits the effective classification performance for rare species. Tabak et al. (2020) suggest increasing the diversity of the camera trap sites and therefore the backgrounds on which to train the models (Tabak et al., 2020).

Mendoza et al. (2011) proposed a new method to help reduce individual identification bias. For this, an online web interface was constructed that allows first to classify all the pictures captured by camera traps into time-related clusters and then allows the classifiers to independently name the target species of bobcats simultaneously, in a mutually blind procedure. The picture identification tool significantly decreased the differences between the classifiers as shown by the Adjusted Rand Index (ARI) (Mendoza et al., 2011).

Delisle et al. (2021) predict that the number of studies that set out to examine unmarked populations will grow, these studies will highly rely on modern technology and relatively newly developed methods, especially for population abundance estimations (Delisle et al., 2021). Delisle et al. (2021) suggest that improved accessibility of software and continued methodological refinement should result in better decision-making and greater adoption by future camera trap studies focused on abundance estimation (Delisle et al., 2021).

Acknowledgements

The authors would like to thank the Tempus Public Foundation for providing the Stipendium Hungaricum scholarship and we extend our great thanks to The Doctoral School of Animal Biotechnology and Animal Science of the Hungarian University of Agriculture and Life Sciences (MATE).

References

Avrin, A. C., Pekins, C. E., Sperry, J. H., Wolff, P. J., & Allen, M. L. (2021). Efficacy of attractants for detecting eastern spotted skunks: an experimental approach. *Wildlife Biology* **2021**(4), wlb.00880. doi: 10.2981/wlb.00880

Balme, G. A., Hunter, L. T. B., & Slotow, R. (2009). Evaluating Methods for Counting Cryptic Carnivores. *The Journal of Wildlife Management* **73**(3), 433-441. doi: 10.2193/2007-368

Balme, G. A., Slotow, R., & Hunter, L. T. (2009). Impact of conservation interventions on the dynamics and persistence of a persecuted leopard (*Panthera pardus*) population. *Biological Conservation* **142**(11), 2681-2690. doi: 10.1016/j.biocon.2009.06.020

Barcelos, D. C., Alvarenga, G. C., Gräbin, D. M., Baccaro, F., & Ramalho, E. E. (2023). Divergent effects of lure on multi-species camera-trap detections and quality of photos. *Journal for Nature Conservation* **71**(1), 126317. doi: 10.1016/j.jnc.2022.126317

Beirne, C., Sun, C., Tattersall, E. R., Burgar, J. M., Fisher, J. T., & Burton, A. C. (2021). Multispecies modelling reveals potential for habitat restoration to re-establish boreal vertebrate community dynamics. *Journal of Applied Ecology* **58**(12), 2821-2832. doi: 10.1111/1365-2664.14020

Bernard, H., Liew, N. Y. S., Wilson, A., Tangah, J., Tuuga, A., & Matsuda, I. (2022). Inventorying terrestrial mammal species in mixed-mangrove forest of the Lower Kinabatangan, Sabah, Borneo, Malaysia, with special reference to a new locality record of otter civet, *Cynogale bennettii*. *Mammal Research* **67**(1), 31-38. doi: 10.1007/s13364-021-00611-5

Bessone, M., Kühl, H. S., Hohmann, G., Herbing, I., N'Goran, K. P., Asanzi, P., ... Fruth, B. (2020). Drawn out of the shadows: Surveying secretive forest species with camera trap distance sampling. *Journal of Applied Ecology* **57**(5), 963-974. doi: 10.1111/1365-2664.13602

Borah, J., Sharma, T., Das, D., Rabha, N., Kakati, N., Basumatary, A., ... Vattakaven, J. (2014). Abundance and density estimates for common leopard *Panthera pardus* and clouded leopard *Neofelis nebulosa* in Manas National Park, Assam, India. *Oryx* **48**(1), 149-155. doi: 10.1017/s0030605312000373

Burgar, J. M., Burton, A. C., & Fisher, J. T. (2019). The importance of considering multiple interacting species for conservation of species at risk. *Conservation Biology* **33**(3), 709-715. doi: 10.1111/cobi.13233

Burton, A. C., Sam, M. K., Balangtaa, C., & Brashares, J. S. (2012). Hierarchical Multi-Species Modeling of Carnivore Responses to Hunting, Habitat and Prey in a West African Protected Area. *PLoS ONE* **7**(5), e38007. doi: 10.1371/journal.pone.0038007

Buyaskas, M., Evans, B. E., & Mortelliti, A. (2020). Assessing the effectiveness of attractants to increase camera trap detections of North American mammals. *Mammalian Biology* **100**(1), 91-100. doi: 10.1007/s42991-020-00011-3

Cappelle, N., Després-Einspenner, M.-L., Howe, E. J., Boesch, C., & Kühl, H. S. (2019). Validating camera trap distance sampling for chimpanzees. *American Journal of Primatology* **81**(3), e22962. doi: 10.1002/ajp.22962

Comer, S., Speldewinde, P., Tiller, C., Clausen, L., Pinder, J., Cowen, S., & Algar, D. (2018). Evaluating the efficacy of a landscape scale feral cat control program using camera traps and occupancy models. *Scientific Reports* **8**(1), 5335. doi: 10.1038/s41598-018-23495-z

Delisle, Z. J., Flaherty, E. A., Nobbe, M. R., Wzientek, C. M., & Swihart, R. K. (2021). Next-Generation Camera Trapping: Systematic Review of Historic Trends Suggests Keys to Expanded Research Applications in Ecology and Conservation. *Frontiers in Ecology and Evolution* **9**(1), 617996. doi: 10.3389/fevo.2021.617996

Dri, G. F., Blomberg, E. J., Hunter, M. L., Vashon, J. H., & Mortelliti, A. (2022). Developing cost-effective monitoring protocols for track-surveys: An empirical assessment using a Canada lynx *Lynx canadensis* dataset spanning 16 years. *Biological Conservation* **276**(1), 109793. doi:

10.1016/j.biocon.2022.109793

Engeman, R. M., & Witmer, G. W. (2000). IPM strategies: indexing difficult to monitor populations of pest species. In T. Salmon & A. Crabb (Eds.), *Proceedings of the vertebrate pest conference* (Vol. 19, pp. 183–189).

Farris, Z. J., Karpanty, S. M., Ratelolahy, F., & Kelly, M. J. (2014). Predator–Primate Distribution, Activity, and Co-occurrence in Relation to Habitat and Human Activity Across Fragmented and Contiguous Forests in Northeastern Madagascar. *International Journal of Primatology* **35**(5), 859–880. doi: 10.1007/s10764-014-9786-0

Foresman, K. R., & Pearson, D. E. (1998). Comparison of Proposed Survey Procedures for Detection of Forest Carnivores. *The Journal of Wildlife Management* **62**(4), 1217–1226. doi: 10.2307/3801985

Glover-Kapfer, P., Soto-Navarro, C. A., & Wearn, O. R. (2019). Camera-trapping version 3.0: current constraints and future priorities for development. *Remote Sensing in Ecology and Conservation* **5**(3), 209–223. doi: 10.1002/rse2.106

Gompper, M. E., Kays, R. W., Ray, J. C., LaPoint, S. D., Bogan, D. A., & Cryan, J. R. (2006). A comparison of noninvasive techniques to survey carnivore communities in northeastern North America. *Wildlife Society Bulletin* **34**(4), 1142–1151. doi: 10.2193/0091-7648(2006)34[1142:ACONTT]2.0.CO;2

Howe, E. J., Buckland, S. T., Després-Einspenner, M.-L., & Köhl, H. S. (2017). Distance sampling with camera traps. *Methods in Ecology and Evolution* **8**(11), 1558–1565. doi: 10.1111/2041-210X.12790

Jordan, M. J., Barrett, R. H., & Purcell, K. L. (2011). Camera trapping estimates of density and survival of fishers *martes pennanti*. *Wildlife Biology* **17**(3), 266–276. doi: 10.2981/09-091

Joubert, C. J., Tarugara, A., Clegg, B. W., Gandiwa, E., & Muposhi, V. K. (2020). A baited-camera trapping method for estimating the size and sex structure of African leopard (*Panthera pardus*) populations. *MethodsX* **7**(1), 101042. doi: 10.1016/j.mex.2020.101042

Kämmerle, J.-L., Ritchie, E. G., & Storch, I. (2019). Restricted-area culls and red fox abundance: Are effects a matter of time and place? *Conservation Science and Practice* **1**(11), e115. doi: 10.1111/csp2.115

Kelly, M. J. (2001). Computer-aided photograph matching in studies using individual identification: An example from Serengeti cheetahs. *Journal of Mammalogy* **82**(2), 440–449. doi: 10.1644/1545-1542(2001)082<0440:capmis>2.0.co;2

Kelly, M. J., Betsch, J., Wultsch, C., Mesa, B., & Mills, L. S. (2012). Noninvasive sampling for carnivores. In L. Boitani & R. A. Powell (Eds.), *Carnivore ecology and conservation: a handbook of techniques* (p. 47–69). Oxford University Press New York.

Kluever, B. M., Gese, E. M., Dempsey, S. J., & Knight, R. N. (2013). A comparison of methods for monitoring kit foxes at den sites. *Wildlife Society Bulletin* **37**(2), 439–443. doi: 10.1002/wsb.261

Larrucea, E. S., Brussard, P. F., Jaeger, M. M., & Barrett, R. H. (2007). Cameras, Coyotes, and the Assumption of Equal Detectability. *The Journal of Wildlife Management* **71**(5), 1682–1689. doi: 10.2193/2006-407

Larrucea, E. S., Serra, G., Jaeger, M. M., & Barrett, R. H. (2007). Censusing bobcats using remote cameras. *Western North American Naturalist* **67**(4), 538–548. doi: 10.3398/1527-0904(2007)67[538:cburc]2.0.co;2

Lazenby, B. T., Mooney, N. J., & Dickman, C. R. (2015). Detecting species interactions using remote cameras: effects on small mammals of predators, conspecifics, and climate. *Ecosphere* **6**(12), 1–18. doi: 10.1890/ES14-00522.1

Macdonald, D. W., Chiaverini, L., Bothwell, H. M., Kaszta, Ž., Ash, E., Bolongon, G., ... Cushman, S. A. (2020). Predicting biodiversity richness in rapidly changing landscapes: climate,

low human pressure or protection as salvation? *Biodiversity and Conservation* **29**(14), 4035-4057. doi: 10.1007/s10531-020-02062-x

Meek, P., Ballard, G., Fleming, P., & Falzon, G. (2016). Are we getting the full picture? Animal responses to camera traps and implications for predator studies. *Ecology and Evolution* **6**(10), 3216-3225. doi: 10.1002/ece3.2111

Mendoza, E., Martineau, P. R., Brenner, E., & Dirzo, R. (2011). A novel method to improve individual animal identification based on camera-trapping data. *The Journal of Wildlife Management* **75**(4), 973-979. doi: 10.1002/jwmg.120

Miyamoto, K., Squires, T. E., & Araki, H. (2018). Experimental evaluation of predation of stocked salmon by riparian wildlife: effects of prey size and predator behaviours. *Marine and Freshwater Research* **69**(3), 446-454. doi: 10.1071/MF17215

Muench, C., & Martínez-Ramos, M. (2016). Can Community-Protected Areas Conserve Biodiversity in Human-Modified Tropical Landscapes? The Case of Terrestrial Mammals in Southern Mexico. *Tropical Conservation Science* **9**(1), 178–202. doi: 10.1177/194008291600900110

Negrões, N., Sarmiento, P., Cruz, J., Eira, C., Revilla, E., Fonseca, C., ... Silveira, L. (2010). Use of camera-trapping to estimate puma density and influencing factors in central Brazil. *The Journal of Wildlife Management* **74**(6), 1195-1203. doi: 10.1111/j.1937-2817.2010.tb01240.x

Nekaris, K. A. I., Handby, V., Campera, M., Birot, H., Hedger, K., Eaton, J., & Imron, M. A. (2020). Implementing and Monitoring the Use of Artificial Canopy Bridges by Mammals and Birds in an Indonesian Agroforestry Environment. *Diversity* **12**(10), 399. doi: 10.3390/d12100399

Nichols, J. D., & Williams, B. K. (2006). Monitoring for conservation. *Trends in Ecology & Evolution* **21**(12), 668-673. doi: 10.1016/j.tree.2006.08.007

O'Brien, T. G., & Kinnaird, M. F. (2011). Density estimation of sympatric carnivores using spatially explicit capture–recapture methods and standard trapping grid. *Ecological Applications* **21**(8), 2908-2916. doi: 10.1890/10-2284.1

O'Connell, A. F., Nichols, J. D., & Karanth, K. U. (Eds.). (2011). *Camera traps in animal ecology: Methods and analyses*. Springer Japan. doi: 10.1007/978-4-431-99495-4

Palencia, P., Rowcliffe, J. M., Vicente, J., & Acevedo, P. (2021). Assessing the camera trap methodologies used to estimate density of unmarked populations. *Journal of Applied Ecology* **58**(8), 1583-1592. doi: 10.1111/1365-2664.13913

Palmer, R., Anderson, H., Richards, B., Craig, M. D., & Gibson, L. (2021). Does aerial baiting for controlling feral cats in a heterogeneous landscape confer benefits to a threatened native meso-predator? *PLOS ONE* **16**(5), e0251304. doi: 10.1371/journal.pone.0251304

Pirie, T. J., Thomas, R. L., & Fellowes, M. D. E. (2016). Limitations to recording larger mammalian predators in savannah using camera traps and spoor. *Wildlife Biology* **22**(1), wlb.00855. doi: 10.2981/wlb.00129

Rogan, M. S., Distiller, G., Balme, G. A., Pitman, R. T., Mann, G. K. H., Dubay, S. M., ... O'Riain, M. J. (2022). Troubled spots: Human impacts constrain the density of an apex predator inside protected areas. *Ecological Applications* **32**(4), e2551. doi: 10.1002/eap.2551

Selonen, V., Banks, P., Tobajas, J., & Laaksonen, T. (2022). Protecting prey by deceiving predators: A field experiment testing chemical camouflage and conditioned food aversion. *Biological Conservation* **275**(1), 109749. doi: 10.1016/j.biocon.2022.109749

Shamoon, H., Saltz, D., & Dayan, T. (2017). Fine-scale temporal and spatial population fluctuations of medium sized carnivores in a Mediterranean agricultural matrix. *Landscape Ecology* **32**(6), 1243-1256. doi: 10.1007/s10980-017-0517-8

Silveira, L., Jácomo, A. T., & Diniz-Filho, J. A. F. (2003). Camera trap, line transect census and track surveys: a comparative evaluation. *Biological Conservation* **114**(3), 351-355. doi: 10.1016/S0006-3207(03)00063-6

Sollmann, R., Mohamed, A., Samejima, H., & Wilting, A. (2013). Risky business or simple

solution – Relative abundance indices from camera-trapping. *Biological Conservation* **159**(1), 405-412. doi: 10.1016/j.biocon.2012.12.025

Srbek-Araujo, A. C., Da Cunha, C. J., Roper, J. J., et al. (2017). Post-dispersal seed predation by Atlantic Forest squirrels monitoring lowland tapir latrines. *Tropical Ecology* **58**(3), 673-678.

Steinbeiser, C. M., Kioko, J., Maresi, A., Kaitilia, R., & Kiffner, C. (2019). Relative abundance and activity patterns explain method-related differences in mammalian species richness estimates. *Journal of Mammalogy* **100**(1), 192–201. doi: 10.1093/jmammal/gyy175

Stobo-Wilson, A. M., Brandle, R., Johnson, C. N., & Jones, M. E. (2020). Management of invasive mesopredators in the Flinders Ranges, South Australia: effectiveness and implications. *Wildlife Research* **47**(8), 720. doi: 10.1071/wr19237

Strampelli, P., Andresen, L., Everatt, K. T., Somers, M. J., & Rowcliffe, J. M. (2020). Leopard *Panthera pardus* density in southern Mozambique: evidence from spatially explicit capture–recapture in Xonghile Game Reserve. *Oryx* **54**(3), 405–411. doi: 10.1017/S0030605318000121

Sunarto, S., Kelly, M. J., Parakkasi, K., & Hutajulu, M. B. (2015). Cat coexistence in central Sumatra: ecological characteristics, spatial and temporal overlap, and implications for management. *Journal of Zoology* **296**(2), 104-115. doi: 10.1111/jzo.12218

Tabak, M. A., Norouzzadeh, M. S., Wolfson, D. W., Newton, E. J., Boughton, R. K., Ivan, J. S., ... Miller, R. S. (2020). Improving the accessibility and transferability of machine learning algorithms for identification of animals in camera trap images: MLWIC2. *Ecology and Evolution* **10**(19), 10374-10383. doi: 10.1002/ece3.6692

Thorn, M., Scott, D. M., Green, M., Bateman, P. W., & Cameron, E. Z. (2009). Estimating Brown Hyaena Occupancy Using Baited Camera Traps. *South African Journal of Wildlife Research* **39**(1), 1–10. doi: 10.3957/056.039.0101

Thornton, D., Scully, A., King, T., Fisher, S., Fitkin, S., & Rohrer, J. (2018). Hunting associations of American badgers (*Taxidea taxus*) and coyotes (*Canis latrans*) revealed by camera trapping. *Canadian Journal of Zoology* **96**(7), 769–773. doi: 10.1139/cjz-2017-0234

Trolliet, F., Vermeulen, C., Huynen, M.-C., & Hambuckers, A. (2014). Use of camera traps for wildlife studies: a review. *Biotechnologie, Agronomie, Société et Environnement* **18**(3), 446-454.

Windell, R. M., Bailey, L. L., Young, J. K., Livieri, T. M., Eads, D. A., & Breck, S. W. (2022). Improving evaluation of nonlethal tools for carnivore management and conservation: evaluating fladry to protect an endangered species from a generalist mesocarnivore. *Animal Conservation* **25**(1), 125-136. doi: 10.1111/acv.12726

Xiao, W., Feng, L., Mou, P., Miquelle, D. G., Hebblewhite, M., Goldberg, J. F., ... Ge, J. (2016). Estimating abundance and density of Amur tigers along the Sino–Russian border. *Integrative Zoology* **11**(4), 322-332. doi: 10.1111/1749-4877.12210

Yoshizaki, J., Pollock, K. H., Brownie, C., & Webster, R. A. (2009). Modeling misidentification errors in capture–recapture studies using photographic identification of evolving marks. *Ecology* **90**(1), 3-9. doi: 10.1890/08-0304.1